Partial Quantile Tensor Regression

Dayu Sun, Limin Peng, Zhiping Qiu, Ying Guo, Amita Manatunga Journal of the American Statistical Association

Emory University & Fujian Normal University

September 22, 2025

Classical Tensor Regression

- ► Inputs:
 - ► Tensor covariate $\mathcal{X} \in \mathbb{R}^{p_1 \times \cdots \times p_m}$ (e.g., images, connectivity matrices);
 - Vector covariate $\mathbf{z} \in \mathbb{R}^{p_z}$ (confounders like age, gender).
- ▶ **Output**: Scalar response $y \in \mathbb{R}$ (e.g., disease severity, test score).
- ▶ Model Assumption (conditional mean):

$$\mathbb{E}[y \mid \mathcal{X}, \mathbf{z}] = \alpha + \boldsymbol{\gamma}^{\top} \mathbf{z} + \langle \mathcal{B}, \mathcal{X} \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the tensor Frobenius inner product (Frobenius 内积).

► Training Objective (least squares):

$$\min_{\alpha, \gamma, \mathcal{B}} \frac{1}{n} \sum_{i=1}^{n} \left(y_i - \alpha - \gamma^{\top} \mathbf{z}_i - \langle \mathcal{B}, \mathcal{X}_i \rangle \right)^2.$$

► **Limitation**: Only captures mean effects, unable to describe distributional heterogeneity (分布异质性).

Quantile and Conditional Quantile


Quantile:

$$Q_{\mathbf{y}}(\tau) = \inf\{ \mathbf{q}: \ \Pr(\mathbf{y} \leq \mathbf{q}) \geq \tau \}, \quad 0 < \tau < 1.$$

Find a point q such that the probability of y being to its left is at least τ .

Conditional Quantile (given tensor \mathcal{X}):

$$Q_{y}(\tau \mid \mathcal{X}) = \inf\{ q : \Pr(y \le q \mid \mathcal{X}) \ge \tau \}.$$



Quantile Tensor Regression (QTR)

- ► Inputs:
 - ▶ Tensor covariate $\mathcal{X} \in \mathbb{R}^{p_1 \times \cdots \times p_m}$;
 - ▶ Vector covariate $\mathbf{z} \in \mathbb{R}^{p_z}$.
- ▶ **Output**: Scalar response $y \in \mathbb{R}$.
- Model Assumption (conditional quantile):

$$Q_{y}(\tau \mid \mathcal{X}, \mathbf{z}) = \alpha(\tau) + \boldsymbol{\gamma}(\tau)^{\top} \mathbf{z} + \langle \mathcal{B}(\tau), \mathcal{X} \rangle.$$

► Training Objective (minimize quantile loss):

$$\min_{\alpha(\tau), \gamma(\tau), \mathcal{B}(\tau)} \frac{1}{n} \sum_{i=1}^{n} \rho_{\tau} \Big(y_i - \alpha(\tau) - \gamma(\tau)^{\top} \mathbf{z}_i - \langle \mathcal{B}(\tau), \mathcal{X}_i \rangle \Big).$$

Quantile Loss Function

▶ Quantile Loss Function:

$$\rho_{\tau}(w) = w\{\tau - \mathbb{I}(w < 0)\} = \begin{cases} \tau \ w, & w \ge 0, \\ (\tau - 1)w, & w < 0. \end{cases}$$

► Intuition:

- When $\tau = 0.5 \rightarrow$ Median regression (symmetric penalty, robust to outliers).
- ▶ When $\tau = 0.9 \rightarrow$ Focus more on right tail; $\tau = 0.1 \rightarrow$ left tail.

Advantages:

- Captures effects at different quantiles, robust to outliers and extremes.
- ▶ Different τ address different scientific questions, not just tuning for best performance.

Limitations:

- ▶ Coefficient tensor $\mathcal{B}(\tau)$ is high-dimensional, requires low-rank decomposition (CP/Tucker).
- Optimization requires alternating updates, computationally expensive and slow convergence.

Partial Quantile Tensor Regression (PQTR)

► Model Assumption (conditional quantile):

$$Q_{y}(\tau \mid \mathcal{X}, \mathbf{z}) = \alpha(\tau) + \boldsymbol{\gamma}(\tau)^{\top} \mathbf{z} + \langle \mathcal{B}(\tau), \mathcal{X} \rangle.$$

- ▶ Coefficient tensor $\mathcal{B}(\tau)$ is high-dimensional, difficult to estimate directly.
- ▶ PQTR Idea: Decompose $\mathcal{B}(\tau)$ in Tucker form:

$$\mathcal{B}(\tau) \approx \langle \langle \mathcal{D}(\tau); W_1, W_2, \dots, W_m \rangle \rangle$$

where $W_k \in \mathbb{R}^{p_k \times d_k}$ are factor matrices.

From Objective Function to Quantile Partial Tensor Covariance

Quantile Tensor Regression Objective:

$$\min_{\alpha(\tau), \gamma(\tau), \mathcal{B}(\tau)} \frac{1}{n} \sum_{i=1}^{n} \rho_{\tau} \Big(y_{i} - \alpha(\tau) - \gamma(\tau)^{\top} z_{i} - \langle \mathcal{B}(\tau), \mathcal{X}_{i} \rangle \Big),$$
 where $\rho_{\tau}(w) = w \{ \tau - \mathbb{I}(w < 0) \}.$

Gradient Signal (quantile residual score):

$$R(\tau) = \rho_{\tau} \left(\mathbf{y} - \alpha_{\mathbf{y}|\mathbf{z}}(\tau) - \boldsymbol{\gamma}_{\mathbf{y}|\mathbf{z}}(\tau)^{\top} \mathbf{z} \right), \quad \rho_{\tau}(\mathbf{w}) = \tau - \mathbb{I}(\mathbf{w} < 0).$$

- ▶ Role: Judges if y is above/below quantile τ , as supervision signal.
- ▶ Define Quantile Partial Tensor Covariance:

$$C(\tau) = \mathbb{E}[R(\tau) \mathcal{X}].$$

Analogous to cov(y, X) in PLS, but using quantile residuals.



From Direct Estimation of $\mathcal{B}(\tau)$ to Regression on Low-Dim Core

Original Objective:

$$\min_{\alpha, \gamma, \mathcal{B}(\tau)} \frac{1}{n} \sum_{i=1}^{n} \rho_{\tau} \Big(y_i - \alpha - \gamma^{\top} z_i - \langle \mathcal{B}(\tau), \mathcal{X}_i \rangle \Big).$$

Tucker Reparameterization:

$$\mathcal{B}(\tau) = \langle \langle \mathcal{D}(\tau); W_1, \dots, W_m \rangle \rangle.$$

Key Identity (inner product invariance under multi-mode projection):

$$\langle \mathcal{B}(\tau), \mathcal{X}_i \rangle = \langle \mathcal{D}(\tau), \underbrace{\mathcal{X}_i \times_1 W_1^\top \cdots \times_m W_m^\top}_{\mathcal{T}_i(\tau)} \rangle.$$

 \triangleright Equivalent Objective (estimate W_k first, then estimate low-dim coefficients):

$$\min_{\alpha, \gamma, \mathcal{D}(\tau)} \frac{1}{n} \sum_{i=1}^{n} \rho_{\tau} \Big(y_{i} - \alpha - \gamma^{\top} z_{i} - \langle \mathcal{D}(\tau), \mathcal{T}_{i}(\tau) \rangle \Big).$$



Estimating Factor Matrices: From $C(\tau)$ to W_k

- Take mode-k unfolding of $C(\tau)$: $C^{(k)}(\tau) \in \mathbb{R}^{p_k \times (p_1 \cdots p_{k-1} p_{k+1} \cdots p_m)}$.
- ► Construct quadratic matrix: $M^{(k)}(\tau) = C^{(k)}(\tau)C^{(k)}(\tau)^{\top}$.
- ► Eigendecomposition: $M^{(k)}(\tau) = U^{(k)} \Lambda^{(k)} U^{(k)\top}$.
- ► Take top d_k eigenvectors: $W_k = U_{[:,1:d_k]}^{(k)}$.

Low-Dim Representation (Tucker Multi-Mode Projection)

Obtain core tensor:

$$\mathcal{T}(\tau) = \mathcal{X} \times_1 W_1^{\top} \times_2 W_2^{\top} \cdots \times_m W_m^{\top},$$

where \times_k is mode-k multiplication.

$$\mathcal{T}(\tau) \in \mathbb{R}^{d_1 \times \cdots \times d_m}$$
.

Perform low-dim quantile regression on $\mathcal{T}(\tau)$.



Theoretical Result 1: Key Identify & Identifiability

Theorem (Pseudo-linearity Identity)

Under mild regularity conditions,

$$\operatorname{vec}(\mathcal{C}(\tau)) = V(\tau)\operatorname{vec}(\mathcal{B}(\tau)), \quad V(\tau) \succ 0.$$

- ▶ Implication 1: $C(\tau) = 0 \iff B(\tau) = 0$ (identifiability).
- ▶ **Implication 2:** The eigenspace of $C(\tau)$ contains the directions of $B(\tau)$.
- ▶ Consequence: One can estimate factor matrices W_k from $C(\tau)$ instead of directly from $B(\tau)$.

Theoretical Result 2: Consistency of PQTR

Theorem (Consistency)

Suppose the true coefficient tensor $\mathcal{B}(\tau)$ satisfies the envelope structure, and the true envelope dimensions are $\{d_k\}$. Then the PQTR estimator

$$\hat{\theta}(\tau) = (\hat{\alpha}, \, \hat{\gamma}, \, \widehat{\mathcal{B}}(\tau))$$

is \sqrt{n} -consistent.

- ► Implication: PQTR achieves the same statistical efficiency as classical quantile regression, even in high dimensions.
- ▶ **Interpretation:** Once the projection dimensions $\{d_k\}$ are chosen correctly, no information is lost —redundancy is removed.

Simulation Setup: Heterogeneous Tri-Square-Shaped Tensor Coefficients

Data Generation:

- ► Response: $Y_i = \gamma^\top Z_i + \langle \mathcal{B}(\xi_i), \mathcal{X}_i \rangle + \Phi^{-1}(\xi_i)$.
- Vector covariate: $\mathbf{Z}_i \in \mathbb{R}^{p_z}$.
- ▶ Tensor covariate: $\mathcal{X}_i \in \mathbb{R}^{50 \times 50}$, covariance $\Sigma_2 \otimes \Sigma_1$.
- Coefficient tensor: $\mathcal{B}(\tau)$ is heterogeneous step function: $\mathcal{B}(\tau) = \mathcal{B}_1 \mathbb{I}(0 < \tau < 0.35) + \mathcal{B}_2 \mathbb{I}(0.35 \le \tau < 0.65) + \mathcal{B}_3 \mathbb{I}(0.65 \le \tau < 1)$, with different ranks (e.g., 1,2,3) and bases (tri-square-shaped).

Compared Methods:

- PQTR variants: Fix (oracle ranks), ER (eigenvalue ratio for d_k), CV (5-fold cross-validation for d_k).
- Benchmarks: CP (CP decomposition QR, with fused penalty), TK (Tucker decomposition QR, with lasso), PCA (vectorize then PCA + QR).

Simulation Results




Fig 1. Empirical averages for heterogeneous tri-square-shaped tensor coefficients (step functions of τ) with $\tau=0.1,0.25,0.5,0.75,0.9$, under envelope or diagonal covariate covariance.

Real Data Application Setup: PTSD Neuroimaging Study

Data Source:

- ▶ 98 female subjects with PTSD (post-traumatic stress disorder).
- Response: PTSD symptom severity score (PSS).
- ► Tensor covariate: 279 × 279 functional connectivity matrix from fMRI (resting-state functional magnetic resonance imaging), based on Power et al. (2011) brain parcellation template (脑区划分模板).
- Vector covariate: Age (as confounder).
- Brain networks: Auditory (Aud), Somatomotor Dorsal (SMd), Cingulo Opercular (CO), Default Mode (DMN), Fronto Parietal (FP), Somatomotor Lateral (SMI), Visual (Vis), Salience (Sal).

► Methods Applied:

- ▶ PQTR with ER (eigenvalue ratio) for selecting reduced dimensions d_k (typically 1 or 2 depending on τ).
- ▶ Benchmark: TEPLS (tensor envelope partial least squares, mean-based regression).
- ► **Goal**: Demonstrate PQTR's practical utility in revealing neurobiologically meaningful and interpretable results compared to mean regression.

Real Data Results: Heat Map of Standardized $\hat{\mathcal{B}}(au)$




Fig2. Heat map of standardized $\hat{\mathcal{B}}(\tau)$ by ER (thresholded at cutoff 5), and by TEPLS . Gray lines separate brain functional networks.